

This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo, copyediting, typesetting, and review of the resulting proof before it is published in its final form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Accepted Manuscript

Title: *Helleborus sp.* an ethnopharmacological and toxicological review

Authors: Andzela Brajanovska, Biljana Bauer*

Faculty of Pharmacy, Ss. Cyril and Methodius University, Majka Tereza 47, 1000 Skopje, Republic of Macedonia



DOI:

Received date: September 2018

Accepted date: November 2018

UDC: 615.322:528.675.1

Type of papers: *Review paper*

Please cite this article as:

***Helleborus sp.* an ethnopharmacological and toxicological review**

Andzela Brajanovska, Biljana Bauer*

*Faculty of Pharmacy, Ss. Cyril and Methodius University, Majka Tereza 47,
1000 Skopje, Republic of Macedonia*

Abstract

Helleborus sp. are healing plants from the family *Ranunculaceae*. The aim of this paper is to consider ethnopharmacology and the toxicity of the hellebores which is focused on a detailed research of the value of phytochemicals, as pharmacological attributes of phytomedicine herbs. The data is based on active components. Also, this paper presents the chemical composition of the root in the overall effect of the drug, as well as the importance of the hellebores in pharmacy. It shows a wide range of pharmacological effects such as cardiotoxic, immunostimulative, antibacterial, antitumor, diuretic, and emetic. *Helleborus odorus* Waldst. et Kit. is rich with secondary metabolites such as bufadienolides, flavonoids, phenolic heterozids. With this research it was concluded that these plants have great significance and great application in the formulation of pharmaceutical dosage forms.

Keywords: *Helleborus sp.*, phytomedicine, folk medicinal uses, cardiotoxic, toxicity, hellebore

Introduction

Medicinal herbs have been used for healing from the beginning of the historical development of mankind. When primal people were suffering from some disease, they were directed to nature, seeking help from it. Surrounded by herbs, insects, animals, and hunted by the self-preservation instinct, people used them to fight the disease. Personal, or following the general and animal samples, people made a selection of herbs, although they often paid for it with life. Numerous literary data indicate the early application of hellebores in traditional folk medicine.

Helleborus sp. and its use began in old civilizations, more than 2500 BC, when ancient Greeks used *Helleborus niger* L. (*Helleboros melas*) because of his own strong diuretic, emetic

and narcotic effects, which cures deafness, leprosy, scabies and psychological disorders (Vitalini et al., 2011). This species today dominates the flora of Southern and Eastern Europe. Orators in ancient Greece used infusions prepared from the leaves of *Helleborus cyclophyllus* Boiss. which strengthened their voice. The old people used hellebores to spread their arrows either when they went on hunting or in wars (Tucakov, 1996). Hippocrates (459-377 BC) and Theophrastus (370-287 BC) have data on the functioning of the hellebores at constipation as laxative and emetic. The Roman medical writer Celsius (25 BC to 50 BCE) in the book “*De re medica*” and physician Galen recorded the medicinal effect of the hellebores in reducing pain (besides cannabis, opium), causing sneezing, in mental disorders (psychosis and depression) and epilepsy. The Romans made a diuretic beverage from this plant that helped to remove the toxins from the body (Van Tellinghen, 2007).

In the Italian folk medicine, underground parts of hellebores in the shape of decoct as a cardiotoxic agent and as a remedy for the purpose of improving the digestive tract was used (Coassini and Poldini, 1988). Alcohol extract from the stem, rhizome and root acts directly in the treatment of migraine, and is also used as an anabolic, vasodilator and antiseptic medicine (Tosevski, 1999). Dried rhizome from *Helleborus thibetanus* Franch. Is used in folk medicine for the treatment of cysts and traumatic injuries (Yang et al., 2010).

Regardless of religion and social order, all nations in the world have, in a greater or lesser extent, usage of hellebores for healing of various diseases. In this paper, the action, as well as the active components of the hellebore, *Helleborus odoratus* Waldst. et Kit. (Syn.: *H. cyclophyllus* Boiss.), a herb of the family *Ranunculaceae*, has been taken in considerations, because it can also be a drug or poison for the heart.

Taxonomy

According to the current scientific systematics of the herbarium, the taxonomic species *Helleborus odoratus* Waldst. et Kit. Is from the kingdom (Regnum) *Plantae* s. *Vegetabilia* (most herbs Cormobionta), division *Magnoliophyta* s. *Angiospermae* (flowers Anthopyta), classis dicotyledone *Magnoliopsida* s. *Dicotyledones*, ordo *Ranunculales* s. *Ranales*, familia *Ranunculaceae*, subfamilia *Ranunculoideae*, genus *Helleborus* L., subgenus *Helleborastum* (Gajić, 1992).

Hellebores are found in the forests or beside them and along the pastures up to 1500 meters, very often throughout Republic of Macedonia. Many authors have cited *Helleborus cyclophyllus* Boiss. From Macedonia as a special species (Micevski, 1985).

Phytochemistry

Early in the 19th century many researchers began researching the chemical structure and the effects of many different plant species. First detailed paper on the chemical structure of the genus *Helleborus* L., of the identification, isolation, characteristics and the biological activity of certain ingredients was done by Husemann and Marmé in 1865. Modern analytical methods have made possible to isolate and thoroughly investigate the structure, biogenesis and the role of various ingredients of the hellebores, but this primary and secondary metabolites, has not yet been sufficiently studied on the human organism.

Primary metabolites in the genus *Helleborus* L.

Many authors, beside the pharmacologically active substances from the underground organs of the hellebores, also examined the content of the primary metabolites. Glucose and other carbohydrates occur by photosynthesis process, transformation and polymerization, as well as with glycolysis, there is the formation of fatty acids and lipids, amino acids and peptides.

Carbohydrates

Carbohydrates are universal ingredients of all living organisms and represent the energy source for the cells. In the plant tissue have a material role, they are integral parts metabolites and precursors of secondary metabolites (Kovacević, 2000). The biochemical Bourquelot method of enzymatic hydrolysis of heterosides was performed on previously stabilized rhizomes and roots of *Helleborus odorus* Waldst. et Kit. And *Helleborus atrorubens* Waldst. et Kit. (Stefanović, 1962). For hydrolysis, the follow enzymes were used: invertin (acting on α -glucosides) and emulsine (acting on γ -glucosides). By comparing of the optical activity of the 3temperature and the amount of reductive substances before and after hydrolysis and the calculated enzymolytic reduction index, it was noted that in the underground organs of *Helleborus odorus* Waldst. et Kit. Besides sucrose,

there are both raffinose and starchy or one of them, and in *Helleborus atrorubens* Waldst. et Kit. There is only saccharose. Enzyme preparation obtained from the leaves and from the underground hellebores organs gave negative results. Stefanovic (1962) from 96% alcoholic extract of underground organ of hellebores, by chromatography method on the column of aluminum oxide (Merck) isolate sucrose, thereby was confirmed her presence in the underground organs of *Helleborus odorus* Waldst. et Kit. (in small quantities) and in *Helleborus atrorubens* Waldst. et Kit.(3%), which was previously been determined only by a biochemical method. Then, by chromatography on paper at *Helleborus odorus* Waldst. et Kit. Were identified: raffinose, sucrose, galactose, glucose, fructose, and at *Helleborus atrorubens* Waldst. et Kit. In addition to the above, also lyxose was present. Baytop and Malkoc, 1965 in the infusions and tinctures of underground organs of *Helleborus orientalis* Lam. Find glucose, fructose and sucrose (Bottcher, 1965).

Lipid materials

Fatty substances represent the energy reserve of plant cells, and their quantity is not a type of species, but depends on the conditions of the environment (temperature, humidity, mineral matter, soil) and stages of plant development. Most of the oil of the underground organs of the hellebores comprised the widespread classes of lipid substances: phospholipids, mono-, di- and triglycerides, free fatty acids and non-fatty moieties (sterols, esterified sterol derivatives and hydrocarbons in an amount of 0.03-1.5%). They are in the root and rhizomes accumulated in large quantities (14-18%), which is a rare occurrence in the plant world. In the fatty oils of the underground organs and seeds in the genus *Helleborus* L. the most frequent are the triglyceride fractions and free fatty acids, where the dominance of unsaturated fatty acids is an important feature of lipid biosynthesis in plants of this gender. By qualitative and quantitative analysis of the content of oils matter in the underground organs of *Helleborus atrorubens* Waldst. et Kit. Was found that its content in rhizome and root does not qualitatively distinguish, but rhizome (11.12%) is richer in greasy ingredients than the root (8.08%) (Čučković, 1939).

The content of fatty substances from the rhizome and roots of *Helleborus odorus* Waldst. et Kit. Was examined and a higher amount of oil from the root, than from rhizome was obtained, for each month during the vegetation period (Živanov-Stakić and Mladenović, 1971). Minimum amount of fat in both of the underground organs was in December, rising in January, February and March. The highest value reached in April when leaves are formed, it falls in June when the fruit

mature, and from August to November there is a slight increase. In the isolated oil the refractive index, acidic, saponification, ester and iodine number has been determined and the ratio of free, esterified and unsaturated fatty acids was obtained during the growing period. The content of the non-saponified part is determined with gravimetric method. The lowest value was determined in December (1.1%), and the highest in April (5.6%), while with chromatography on a thin layer β -sitosterol, other free and esterified phytosterols were determined. Serebryakov, 1982 and Anguladze, 1983 found that lipids of the rhizome and roots of *Helleborus abchasicus* A. Braun and *Helleborus caucasicus* A. Braun were characterized with high content of free fatty acids (about 70%) in the period of flowering, where linoleic acid (82.6%) is dominant, and the presence of β -sitosterol has been demonstrated. Oxidation of lipids from the underground organs of *Helleborus caucasicus* A. Braun is going in the direction of creation of keto acids and free fatty acids (Dalakishvili, 1983). Variations in the content of fatty substances and fatty acids in the underground organisms of the genus *Helleborus* L. was examined during the ontogenetic development over two years. Colombo, 1991 has found that the amount of fat in *Helleborus odorus* Waldst. et Kit. And *Helleborus viridis* L. is highest during the rest period (since October until February) characterized by relatively high metabolic activity and small growth, the smallest in the spring when the plant uses lipid growth reserves. In *Helleborus niger* L. higher temperatures have a positive effect on the synthesis and accumulation of lipids, so there is the largest amount in the period of growth and development (maximum in July). The content of fatty substances during the entire vegetation period was higher in 5empera of *Helleborus odorus* Waldst. et Kit. And in *Helleborus viridis* L., and at *Helleborus niger* L. in root.

Amino acids and peptides

The amino acids in the cells and tissues of the plants are free or in oligopeptides and proteins, and enter into the composition of some groups of secondary metabolites (alkaloids, sulfur and cyanogenic heterozyme) (Kovacevic, 2000). The method of two-dimensional chromatography on paper, (Stefanović, 1962), on an alcohol extract from the underground parts of *Helleborus odorus* Waldst. et Kit. Identifies and separates the follow amino acids: asparagine, aspartic and glutamic acid, serine, lysine, arginine, glycine, histidine, threonine, alanine, proline, methionine, valine, leucine and isoleucine, while in the extract from *Helleborus atrorubens* Waldst. et Kit. Not detect amino acid serine.

Milbradt et al. (2003) from the rhizome of *Helleborus purpurascens* Waldst. et Kit. Isolated cystine peptide heletin D, potentially useful in tumor therapy. They found the antimicrobial activity and the cytotoxic effect of heletin D, based on its binding to cell phospholipids causing lysis and cell death. It is possible to produce transgenic plants containing thionine genes responsible for increasing resistance to pathogenic microorganisms, fungi and yeast; while on the other hand, hellebores can be used as antitumor support therapy (Grossarth-Maticek et al., 2001).

Pěňčík (2009) has developed a highly specific analytical protocol for isolation and quantification of indole-3-acetic acid (IAA), which is important for the auxin, and its amino acid conjugates having a storage function or degradation of excessive IAA. By combining of anti-IAA immunoaffinity extraction with sensitive and selective LC-MS / MS analysis, with the application of internal standards, in a small amount of unripe seeds (about 30 mg) of *Hellebous niger* L. the physiological levels of these compounds were found in a range of 7.5 nmol / g of IAA to 0.44 pmol / g of IAA conjugated to alanine. IAA level and its conjugates with Asp, Glu and Leu in the pericarp were significantly lower than those in the seed, while the conjugates with Ala, Gly, Phe and Val were below the limits of detection.

Secondary metabolites of the genus *Helleborus* L.

Secondary metabolites are specific, biologically active ingredients that have therapeutic use. Secondary metabolites are due to the metabolic processes of primary metabolites associated with catabolism and/or transformation of carbohydrate, amino acid or fatty acid molecules.

The first analyze of the chemical composition of different species of *Helleborus* was in 1808 year in alcoholic extract of *Helleborus hiemalis* L. (*Eranthis hiemalis* Salisb.), when the resin a substance called heleborin was isolated (Dilparić-Knežević, 1977). Bastik forty-five years later (1853) from *Helleborus niger* L. isolated heleborin in crystal form. The same author points out that Marmé in the middle of the 19th century (1864) from the rhizome and root of *Helleborus niger* L. managed to isolate heleborin, soluble in alcohol and chloroform, and also isolates the new substance heleborein, which was easily soluble in water, hard in alcohol and insoluble in the ether. With drying and evaporating of the aqueous solution transparent yellow resin was obtained. A year later, Husemann and Marmé (1865) published the results of the chemical analyzes of both

substances isolated from the underground organs from *Helleborus niger* L., *Helleborus viridis* L. and *Helleborus foetidus* L. which point out on their glycosidic nature. Heleborin ($C_{36}H_{42}O_6$) by hydrolysis was split to aglycol heleborosin ($C_{30}H_{38}O_4$) and one glucose molecule, and heleborein ($C_{26}H_{44}O_{15}$) to aglycone heleborethine ($C_{14}H_{20}O_3$) and two glucose molecules. They found that heleborin demonstrated narcotic and hemolytic action and is supposed that by composition it was saponin, while heleborein exhibited cardiac function similar to that of digitalis.

Helebrin (Fig. 1) (Karrer, 1943) is the first isolated compound of many others from the underground parts of *Helleborus niger* L. from different habitats in Switzerland, Germany, France, Austria, Hungary and Romania. By extraction with Soxhlet, 7emperatu ($C_{36}H_{52}O_{15}$) was isolated as the only substance with glycosidic character (7emperature7n – glukoraminoside). It is believed that 7emperatu is the major cardiac compound of the genus *Helleborus* L. Helebrin is separated in the form of colorless non – hygroscopic crystal, which in water and ethyl alcohol is difficult to dissolve (about 0.45 or 0.2%), slightly easier in methyl alcohol (about 0.6% - 0.7%), even more easily in dilute methyl and ethyl alcohol, and insoluble in chloroform and ether. Helebrin is physiologically very active: 1 g of 7emperatu contains 25×10^5 - 32×10^5 F.D (Frosch-Dosisrogs doses), while the effectiveness is strongest of the cardioactive glycosides convalatoxin 30×10^5 – 35×10^5 F.D./g. Lethal dose of 7emperatu for adult cats at intravenous administration is 0.1 mg/kg. Oral administration of 1 mg is without symptoms, while a 10 mg dose gives a lethal outcome. Comparing these values with the values for heleborein, Karrer (1943) came to the conclusion that 7emperatu is 20-30 times more effective than heleborein (i.v. heleborein LD 1.9 mg/kg, 7emperatu LD 0.1 mg/kg; heleborein LD 300 mg, 7emperatu LD 10 mg). Shmuz determines the structure of this glycoside, by using a strofantobiasis to separate D-glucose from helibrine. Monoside desglucohelebrine was obtained which was further hydrolyzed by diluted hydrochloric acids in acetone to L-rhamnose and highly active helebrigine (Steinergger, 1972).

Pharmacological activity

Depending on the conditions (doses, application modes), the total extract as well as individual fractions of rhizomes and roots of the hellebores, exhibit multiple bioactive effects and can act in several directions. The mechanisms of action in most cases are not sufficiently precisely examined and described or unknown. Identification and characterization of *Helleborus* L. extract

components involved in the induction of certain effects are of primary importance for clarifying the basic mechanisms of their *in vitro* and *in vivo* activity.

Immunostimulative effect

In animal health care, immunomodulatory substances are used, which by different mechanisms, change the level of activity of the components of the immune system. Exogenous immunomodulators stimulate the immune response of the cellular and humoral type (immunostimulators) or reduce unwanted effects of the immune system activity and they encourage mechanisms of tolerance in order to reduce the negative response to their own and allo-antigens (immunosuppressants prevent transplant rejection, allergic and autoimmune reactions). A large number of substances have been identified with immunomodulatory substances from chemical products to products of plant origin, and it is confirmed that the extract of rhizomes and the root of the hellebores acts immunomodulatory (Rosselli et al., 2007).

Cardiotonic effect

Glycosides from *Helleborus odorus* Waldst. et Kit. Have a expressive cardiovascular effect. The effect is somewhat more expensive than that of the digitalis, and worse than the strophanth and less accumulates from the digitalis glycosides. Heleborin is the main active substance in the *Helleborus odorus* Waldst. et Kit. And is the carrier of cardiovascular effect. Heleborein is more effective than k-strophabrizide in intravenous administration, but it is very toxic: the lethal dose (LD) of k-strophantozide is 0.260 mg / kg, while of heleborin is 0.176 mg/kg. *Helleborus odorus* Waldst. et Kit. Have a lot of applications in veterinary medicine. It is prepared as decoct as antiparasitic medicine and as a purgative (Rosselli et al., 2007). In contact with skin and mucous membranes, *Helleborus odorus* Waldst. et Kit. Manifest a very irritating effect, followed by burning and redness, and rarely dermatitis.

Antibacterial effect

Rosselli and coworkers, 2007 analyzed the antibacterial action of the extract of *Helleborus bocconei* ssp. *Intermedius*, and established that it acts inhibitory to *Staphylococcus epidermalis* and *Staphylococcus aureus*. Root of *Helleborus bocconei* Ten. Ssp. *Siculus* contains substances with antibacterial activity, so is justified the usage of this herb in traditional treatment to treat respiratory diseases (Puglisi et al., 2009).

Toxical effect

A large number of plant species (more than 1000) contain components which interfere with the metabolism, or have a direct or indirect toxic effect. However, only a small group of these plants leads to severe poisoning after the introduction of a limited amount of plant material, while most cause poisoning only under certain circumstances (Frohne and Pfänder, 2005).

“Third Defense”, work of the Paracelsus (1493-1541) points out: “What is not poison? All things are poison and nothing is not poisonous. The dose only makes the thing a poison” (Deichmann and al., 1986). A well-known principle “*Dosis sola facit venenum*” is equally important for the toxic as well as for medicinal plants; because it indicates that the degree of toxicity of a particular plant species depends on the dose entered in organism (Mulet, 1991). Toxicity of hellebores depends on the degree of maturity or phenophase (leaf, flowering, fruiting), from plant organism (accumulation of toxic substances takes place in the leaf, fruits, seeds or underground organs) and whether it is used fresh or dried (Spoerke and Smolinske, 1990).

Pre-conditions for poisoning are: contact with the plant, a high enough dose of toxic ingredients to exhibit activity in or on body and defensive defense mechanisms and detoxification processes of the organism. Level poisoning of animals depends on the type of animal, nutrition, health status, age and pregnancy. Young, poorly-nourished and gravid animals are more likely to get sick and more severe clinical form of the poisoning (Frohne and Pfänder, 2005).

In phytotherapy the term “toxicity or toxic plants” can be conditionally used, since many of them are used as medicines in traditional and conventional medicine, sometimes as an antidote. New findings suggest that the hellebores take a significant place in systematic therapy, which is consulted with alternative medicine labeled as homeopathy. Its application stimulates mental and physiological functions in neurological and mental regression, and is also indicated in septicemia, autism, learning difficulties and act in psychosis in young women.

It is thought that Alexander the Great, died as a result of poisoning with this hellebore with which was spiced up the wine he drank (Deichmann, 1986). From previous investigations that were made, it was concluded that the hellebores could also be a poison and a medicine for the heart, because it contains bufadienylid heterozids, which in small therapeutic doses act toxic to the heart (cardiotonic), but because of its low therapeutic width, it can act cardiotonic.

Conclusion

Helleborus sp. Are a medical plants originating from the *Ranunculaceae* family that attracted researchers for its traditional medicinal uses and highly valued bioactive substances. Through this paper, the systematics of the hellebores, as well as the chemical composition, the action and the use of these plants are considered. Because of their toxicity, these plants are strictly limited in medicine as well as for folk purposes because they contain bufadienylid heterozids, which in small therapeutic doses act toxic to the heart (cardiotonic) similar to the digitalis, which this herbs placed in the group of poisonous plants. The medicinal and toxic properties of these plants originate from the secondary metabolites that are biochemical or physiologically active in the body.

References

- Adcock, I.M., Caramori, G., 2001. Cross-talk between pro-inflammatory transcription factors and glucocorticoids. *Immunol. Cell Biol.* 79(4), 376-384.
- Bagrov, A.Y., Shapiro, J.I., Fedorova, O.V., 2009. Endogenous cardiotonic steroids: physiology, pharmacology, and novel therapeutic targets. *Pharmacol. Rev.* 61(1), 9-38.
- Bottcher, H., 1965. *Miracle drugs*, Zora, Zagreb.
- Coassini, L., Poldini, L., 1988. Herbal remedies in the traditional medicine of the Venezia Giulia Region (North East Italy). *J Ethnopharm.* 22(3), 231-79.
- Čučković, D., 1939. Prilog farmakognoškom istraživanju crnocrvenog kukurijeka (*Helleborus atrirubens* Waldst. et Kit.). Doktorska disertacija, Zagreb.

- Dalakishvili, T.M., 1983. Fatty acidic composition of lipids from the roots and rhizomes of *Helleborus caucasicus*. Soobshch Akad Nauk Gruz SSR. 112(3), 565-568.
- Deichmann, W.B., Henschler, D., Holmstedr, B., Keil, G., 1986. What is there that is not a poison? A study of the Third Defence by Paracelsus. Arch. Toxicol. 58, 207-213.
- Dervendzi, V., 1986. Prirodni lekoviti I aromaticni surovini. Nasa knjiga, Skopje.
- Dilparić-Knežević, E., 1977. Bufadienolidi I bazične supstancije u vrstama roda *Helleborus*. Magistarska teza, Zagreb.
- Frohne, D., Pfänder, H.J., 2005. Poisonous Plants, A Handbook for Doctors, Pharmacists, Toxicologists, Biologists and Veterinarians, Second ed. Manson Publishing Ltd.
- Gajić, M., 1992. Fam. *Ranunculaceae* A. L. Juss. In: Sarić RM I ost., Flora Srbije 1, Srpska akademija nauka I umetnosti. Odeljenje prirodno matematičkih nauka. Beograd, 274-280.
- Gligič, V., 1953. Etimološki botanički rečnik. "Veselin Masleša", Sarajevo.
- Goronovic, S.M., Lukic, P., 2001. Farmakognozija. Farmaceutski fakultet Univerziteta u Beograd, Beograd.
- Grossarth-Maticsek, R., Kiene, H., Baumgartner, S.M., Ziegler, R., 2001. Use of Iscador, an extract of European mistletoe (*Viscum album*), in cancer treatment: prospective nonrandomized and randomized matched-pair studies nested within a cohort study. Altern. Ther. Health Med. 7(3), 57-78.
- Hajrulai, Z., 2001. Efektot na ekstraktot na *Helleborus odoratus* vrz proteinskiot status kaj jaginjata. Doktorska disertacija, „Veterinaren fakultet“, Skopje.
- Karrer, W., 1943. Über Hellebrin, ein krystallisiertes Glykosid aus Radix *Hellebori nigri*. Helv. Chim. Acta 26, 1353-1367.
- Kißmer, B., Wichtl, M., 1986. Bufadienolide aus Samen von *Helleborus odoratus*. Planta Med. 2, 152-153.
- Kolesnikov, D.G., Tropp, M.I., 1952. Cardiac glycosides from root of *Helleborus*. Med. Prom. SSSR. 6(5), 17-20.
- Kovacevic, N., 2000. Osnovi Farmakognozije. Licno izdanje, Beograd.
- Micevski, K., 1985. Flora na SR Makedonija. Makedonska akademija na nauki I umetnost, Skopje.
- Mulet, L., 1991. Estudio Ethnobotánico de la Provincia de Castellón, Diputación de Castellón, Castellón, Spain, 596.

- Pěňčík, A., Rolčík, J., Novák, O., Magnus, V., Barták, P., Buchtík, R., Salopek-Sondi, B., Strnad, M., 2009. Isolation of novel indole-3-acetic acid conjugates by immunoaffinity extraction. *Talanta* 80, 651-655. Available at: <https://doi.org/10.1016/j.talanta.2009.07.043>.
- Plowden, C.C., 1970. *A Manual of Plant Names*. George Allen & Unwin, London.
- Puglisi, S., Speciale, A., Acquaviva, R., Ferlito, G., Ragusa, S., De Pasquale, R., Iauk, L., 2009. Antibacterial activity of *Helleborus bocconei* Ten. Subsp. Siculus root extracts. *J. Ethnopharmacol.* 125, 175-177. Available at: <https://doi.org/10.1016/j.jep.2009.06.011>.
- Rosselli, S., Maggio, A., Formisano, C., Napolitano, F., Senatore, F., Spadaro, V., Bruno, M., 2007. Chemical composition and antibacterial activity of extracts of *Helleborus bocconei* Ten. Subsp. Intermedius. *Nat. Prod. Commun.* 2, 675- 679. Available at: <https://doi.org/10.1177/1934578X0700200611>.
- Silic, C., 1987. *Sumske zeljaste biljke*. Svjetlost, Sarajevo.
- Steyn, P.S., Van Heerden, F.R., 1998. Bufadienolides of plant and animal origin. *Nat. Prod. Reports.* 15, 397-413. Available at: <https://doi.org/10.1039/a815397y>.
- Souto, W.M., Mourão, J.S., Barboza, R.R., Alves, R.R., 2011-b. Parallels between zootherapeutic practices in ethnoveterinary and human complementary medicine in northeastern Brazil. *J. Ethnopharmacol.* 134(3), 753-767. Available at: <https://doi.org/10.1016/j.jep.2011.01.041>.
- Spoerke, J.D.G., Smolinske, S.C., 1990. *Toxicity of Houseplants*. CRC Press: Boca Raton, FL, 37-39.
- Sprung, C.L., Caralis, P.V., Marcial, E., Pierce, M., Gelbard, M.A., Long, W.M., Duncan, R.C., Tendler, M.D., Karpf, M., 1984. The effects of high-dose corticosteroids in patient with septic shock. A perspective, controlled study. *N. Engl. J. Med.* 311(18), 1137-1143.
- Stefanović, O., 1962. Ispitivanje glucida *Helleborus*-a biohemijskom metodom. *Acta Pharm. Jug.* 12, 139-146.
- Steinegger, E., Hänsel, R., 1972. *Lehrbuch d. allgemeinen Pharmakognosie*. Springer Verlag, Berlin – Heidelberg – New York.
- Stroman, F., Jakovlev, V., Metzner, P., Roth, E., Thiemer, K., 1984. The pharmacology of a new cardiosteroid, a partially synthetic derivative of the aglycone hellebrigenin (acrihellin). *Arzneimittelforschung.* 34(7), 769-779.

- Tosevski, J., Stojkovski, V., Ulčar, I., 1999. Određivanje na 13temperature koeficient na svinjite so upotreba na ekstrakt od bilkata *Helleborus odorus*. Zbornik na trudovi, XXIV sredba "Fakultet – stopanstvo" '99, Skopje 7, 177- 189.
- Tucakov, J., 1971. Healing with plants – phytotherapy. Culture, Beograd.
- Tucakov, J., 1996. Lečenje biljem. Rad, Beograd.
- Van Tellingen, C., 2007. Pliny's pharmacopoeia of the Roman treatment. Neth. Heart J. 15, 118-120. Available at: <https://doi.org/10.1007/bf03085966>.
- Vitalini, S., Braca, A., Fico, G., 2011. Study on secondary metabolite content of *Helleborus niger* L. Leaves. Fitoterapia 82, 152-154. Available at: <https://doi.org/10.1016/j.fitote.2010.08.012>.
- Wiat, C., 2006. Ethnopharmacology of medicinal plants. Humana Press, New Jersey.
- Wittmann, V., 2008. Glycoproteins: Occurrence and Significance, Glycoscience: Chemistry and Chemical Biology; Fraser-Reid, B.O., Tatsuta, K., Thiem, J., Eds. Springer, Berlin/Heidelberg.
- Yang, J., Zhang, Y., Miao, F., Zhou, L., Sun, W., 2010. Two new bufadienolides from the rhizomes of *Helleborus thibetanus* Franch. Fitoterapia 81, 636-639. Available at: <https://doi.org/10.1016/j.fitote.2010.03.009>.
- Živanov-Stakić, D., Mladenović, M., 1970-a. Novi postupak dobijanja triterpenskikh saponozida biljke *Helleborus odorus*. Acta Pharm. Jug. 20, 183- 186.

Резиме

Етнофармаколошки и токсиколошки преглед на *Helleborus sp.*

Анцела Брајановска, Билјана Бауер*

Фармацевтски факултет, Универзитет „Св. Кирил и Методиј“, Мајка Тереза 47,
1000 Скопје, Република Македонија

Клучни зборови: *Helleborus sp.*, фитомедицина, употреба во народната медицина, кардиотоник, токсичност, кукурек

Helleborus sp. се лековити растенија од фамилијата *Ranunculaceae*. Целта на овој труд е да се разгледа етнофармакологијата и токсичноста на различни видови кукурек која е фокусирана на детални истражувања на вредноста на фитохемикалиите, како фармаколошки атрибути на фитомедицинските билки. Податоците се изложени врз основа на активните компоненти. Исто така во овој труд е изложен хемискиот состав на коренот во целокупното дејство на дрогата како и важноста на различни видови кукурек во фармацијата. Различните видови кукурек покажуваат широк спектар на фармаколошко дејство како што е кардиотонично, имуностимулативно, антибактериско, антитуморно, диуретично и еметично дејство. *Helleborus odorus* Waldst. et Kit. е богат со секундарни метаболити како што се буфаденолиди, флавоноиди, фенолни хетерозиди. Со ова истражување е заклучено дека ова растение има големо значење и голема примена во формулација на фармацевтски дозирани форми.




Fig. 1. Helebrin.

EDITED PROOF